

**This is the author-manuscript version of this work - accessed from  
<http://eprints.qut.edu.au>**

Zhang, Tonghua and Tade, Moses O. and Tian, Yu-Chu (2007) Linear estimate of the number of limit cycles for a class of non-linear systems. *Chaos, Soliton and Fractals* 31(4):pp. 804-810.

**Copyright 2007 Elsevier**

# Linear estimate of the number of limit cycles for a class of non-linear systems<sup>★</sup>

Tonghua Zhang<sup>a,1</sup>, Moses O. Tadé<sup>a</sup>, Yu-Chu Tian<sup>b</sup>

<sup>a</sup>*Department of Chemical Engineering, Curtin University of Technology, GPO Box  
U1987, Perth WA 6845, Australia*

<sup>b</sup>*School of Software Engineering and Data Communications, Faculty of  
Information Technology, Queensland University of Technology, GPO Box 2434,  
Brisbane QLD 4001, Australia*

---

## Abstract

A dynamic system has a finite number of limit cycles. However, finding the upper bound of the number of limit cycles is an open problem for general non-linear dynamical systems. In this paper, we investigated a class of non-linear systems under perturbations. We proved that the upper bound of the number of zeros of the related elliptic integrals of the given system is  $7n + 5$  including multiple zeros, which also gives the upper bound of the number of limit cycles for the given system.

---

## 1 Introduction

Limit cycle behavior has been observed in many application-oriented dynamical systems. Typical examples of limit cycle behavior include feedback systems [19]; industrial optimization [7]; switched fluid networks for communications, computer systems, and flexible manufacturing [13]; continuous stirred-tank reactors in chemical engineering (CSTR) [17]; machine tool chatters [24], and the generalized Rayleigh-Liénard oscillator [16]. Limit cycles are also observed in the Liénard system, an important class of nonlinear dynamical systems in physics and engineering [6,12,14,16]. They play an important role in the dynamic behavior of many practical systems. The following two aspects are of

---

<sup>★</sup> This work is supported in part by Australian Research Council (ARC) under the Discovery Projects scheme (grant ID: DP0559111).

<sup>1</sup> Corresponding author. Tel.: +61-8-9266 7569, fax: +61-8-9266 2681, e-mail address: t.zhang@exchange.curtin.edu.au (T. Zhang).

particular interest for charactering the limit cycles of a system qualitatively and quantitatively:

- (1) How many limit cycles can a system have?
- (2) What is the distribution of the limit cycles?

Research on limit cycle behavior is also important for theoretical advances, and has attracted the interests of many mathematicians ([2,4,5,8–11,15,18,20,21,25]). Consider the following dynamical system

$$\begin{aligned}\dot{x} &= \sum_{i+j=0}^n a_{ij}x^i y^j = P(x, y) \\ \dot{y} &= \sum_{i+j=0}^n b_{ij}x^i y^j = Q(x, y)\end{aligned}\tag{E_n}$$

where  $(x, y) \in \mathbb{R}^2$ ,  $a_{ij}, b_{ij} \in \mathbb{R}$ . The second part of the Hilbert's 16<sup>th</sup> problem asks the uniformly bounded upper bound of the number of limit cycles of system  $(E_n)$ . But this problem has been so difficult that there has been no elegant method to solve it, even for  $n = 2$ . In 1977, Arnold [2] posed a simpler problem, which was called the weakened Hilbert's 16<sup>th</sup> problem or the Hilbert-Arnold problem:

Let  $H$  be a real polynomial with  $\deg H = n$  and  $f$  be a polynomial with  $\deg f = m$ . What is the upper bound of the number of zeros of the following integral?

$$I(h) = \iint_{H \leq h} f dx dy$$

In this paper, we consider a class of non-linear systems of the form

$$\begin{aligned}\dot{x} &= y + \varepsilon P(x, y), \\ \dot{y} &= -(x^5 + bx^3 + x) + \varepsilon Q(x, y)\end{aligned}\tag{1}$$

where  $\varepsilon > 0$  is small, both  $P$  and  $Q$  are polynomials in  $x$  and  $y$ . So, the related elliptic integrals can be expressed as the form

$$I(h) = \oint_{\Gamma_h} Q dx + P dy = \oint_{\Gamma_h} \sum_{k+j \leq n} a_{kj} x^{2k} y^j dx \tag{A}$$

where  $\Gamma_h$  is the level set  $\{H(x, y) = h\}$ . For  $\varepsilon = 0$ , system (1) has a first integral of the form

$$H(x, y) = \frac{y^2 + x^2}{2} + \frac{b}{4}x^4 + \frac{1}{6}x^6 \tag{2}$$

whose level curves have three different types as shown in Fig. 1. They are called the cases of double eight-loops, double-cusps loop, and global center, respectively.




Fig. 1. Level curves of  $H(x, y)$ .

For system (1), Zang et al. proved in [26] that the Hopf cyclicity is two for  $P(x, y) = 0$ ,  $\deg Q(x, y) = 5$  and  $b = -5/2$ . Using the bifurcation theory and qualitative analysis, they also investigated the limit cycles bifurcated from homoclinic and heteroclinic loops; Zhang [23] investigated the Abelian integrals of the system for  $b < -2$ , which is the case (A) in Fig. 1.

The focus of this paper is to investigate the number of zeros of the Abelian integrals of system (1) for  $b \geq -2$ , i.e., the cases (B) and (C) in Fig.1. The paper is organized as follows. We derive the algebraic structure of the Abelian integrals in Section 2. In Section 3, we first prove our result for the case  $b > -2$ ; and then give the result for the case  $b = -2$ . The basic idea is to convert the estimate of the number of zeros of the related elliptic integrals into that of a polynomial in  $h$  through the Picard-Fuchs equations and Riccati equation. Finally, some conclusions are drawn in Section 4.

## 2 Algebraic structure of the Abelian integrals

In this section we will give some relationships that are fundamental for estimating the number of zeros of the Abelian integrals for the case of  $b \geq -2$ . Some results for the case of  $b < -2$  can be found in [23].

Let  $I_{k,j} = \oint_{\Gamma_h} x^{2k} y^j dx$ ,  $I_0 = I_{01}$ ,  $I_1 = I_{11}$ , and  $I_2 = I_{21}$ . Then, the following statement holds.

**Lemma 1** *For  $d \geq 3(k+l=d)$ ,  $I_{kl}$  can be expressed as the linear combination of  $I_{ij}$  (where  $i+j=d-1$ ,  $d-2$ ) and  $hI_{ij}$  (where  $i+j=d-2$ ;  $i=0, 1, 2$ ).*

**PROOF.** It follows from  $H = h$  that

$$\frac{1}{2}I_{k,l+2} + \frac{1}{2}I_{k+1,l} + \frac{b}{4}I_{k+2,l} + \frac{1}{6}I_{k+3,l} = hI_{k,l} \quad (3)$$

and

$$ydy + (x + bx^3 + x^5)dx = 0 \quad (4)$$

Multiplying (4) by  $x^{2k-5}y^l$  and integrating over  $\Gamma_h$  give by part

$$I_{kl} = \frac{2k-5}{l+2}I_{k-3,l+2} - I_{k-2,l} - bI_{k-1,l} \quad (5)$$

Combining Eqs. (3) and (5) gives the following algebraic system

$$\mathbf{A}\mathbf{J} = \mathbf{B}, \quad \mathbf{A} = \begin{pmatrix} \mathbf{A}_5 & \mathbf{0} \\ \mathbf{0} & \mathbf{E}_{d-4} \end{pmatrix} \quad (6)$$

where  $\mathbf{J} = (I_{0,d}, I_{1,d-1}, \dots, I_{d,0})^T$ ,  $\mathbf{E}_{d-4}$  is a unit matrix of order  $d-4$ , and

$$\mathbf{A}_5 = \begin{pmatrix} (6d+2)/d & 0 & b & 0 & 0 \\ 0 & 6d/(d-1) & 0 & b & 0 \\ 0 & 0 & (6d-2)/(d-2) & 0 & b \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}$$

Since  $\det \mathbf{A}_5 \neq 0$  and matrix  $\mathbf{B}$  consists of only integrals  $I_{ij}$  (where  $i+j = d-1, d-2$ ) and  $hI_{ij}$  (where  $i+j = d-2, i=0,1,2$ ), the statement in the lemma is true. This completes the proof.  $\blacksquare$

The straightforward computing by induction yields the following lemma.

**Lemma 2** *For  $n \geq 3$  the Abelian integral  $I(h)$ ,  $h \geq 0$ , can be expressed as the following form*

$$I(h) = \alpha(h)I_0 + \beta(h)I_1 + \gamma(h)I_2 \quad (7)$$

where  $\alpha, \beta, \gamma$  are polynomials in  $h$  with  $\deg \alpha = \left\lfloor \frac{n-1}{2} \right\rfloor$ ,  $\deg \beta = \left\lfloor \frac{n-2}{2} \right\rfloor$ , and  $\deg \gamma = \left\lfloor \frac{n-3}{2} \right\rfloor$ . For  $n = 0$ ,  $\alpha = \beta = \gamma = 0$ ; for  $n = 1$ ,  $\deg \alpha = \beta = \gamma = 0$ ; for  $n = 2$ ,  $\deg \alpha = \deg \beta = \gamma = 0$ .

From [22], we have the following lemma.

**Lemma 3** *The integrals  $I_0, I_1$ , and  $I_2$  satisfy the following Picard-Fuchs equations*

$$\mathbf{T}\mathbf{J} = (3h\mathbf{E} + \mathbf{S})\mathbf{J}' \quad (8)$$

where  $\mathbf{E}$  is the identity matrix of order 3,  $\mathbf{J} = (I_0, I_1, I_2)^T$ , and

$$\mathbf{T} = \begin{pmatrix} 2 & 0 & 0 \\ b/4 & 3 & 0 \\ (4-b^2)/4 & 3b/4 & 4 \end{pmatrix}, \mathbf{S} = \begin{pmatrix} 0 & -1 & -b/4 \\ 0 & b/4 & (b^2-4)/4 \\ 0 & (4-b^2)/4 & (5b-b^3)/4 \end{pmatrix}$$

### 3 Number of zeros of the Abelian integrals

Now we are ready to give a linear estimate of the number of zeros of the abelian integrals. Differentiating both sides of equation (8) with respect to  $h$  gives

$$(\mathbf{T} - 3\mathbf{E})\mathbf{J}' = (3h\mathbf{E} + \mathbf{S})\mathbf{J}'' \quad (9)$$

We now consider different cases for Eq. (9) in the following subsections.

#### 3.1 The case of global center: $b > -2$ and $3b^2 - 16 \neq 0$

For  $3b^2 - 16 \neq 0$ , using the special form of  $\mathbf{T} - 3\mathbf{E}$  yields

$$I_2''(h) = \frac{12bh}{16-3b^2}I_0''(h) + \frac{48h}{16-3b^2}I_1'(h) \quad (10)$$

Let  $Z = \frac{3b}{4}I_1 + I_2$ . Then, Eqs. (8) and (9) show that for  $\mathbf{J}_2 = (I_0, I_1, Z)^T$

$$\mathbf{J}_2 = \mathbf{D}\mathbf{J}'_2, \quad \mathbf{D} = \begin{pmatrix} d_{00} & d_{01} & d_{02} \\ d_{10} & d_{11} & d_{12} \\ d_{20} & d_{21} & d_{22} \end{pmatrix} \quad (11)$$

and

$$G(h)\mathbf{J}_2'' = \mathbf{D}^{-1}(\mathbf{E} - \mathbf{D}')\mathbf{J}'_2 = \mathbf{F} \begin{pmatrix} I'_0 \\ Z' \end{pmatrix}, \quad \mathbf{F} = \begin{pmatrix} f_{00} & f_{02} \\ f_{10} & f_{12} \\ f_{20} & f_{22} \end{pmatrix} \quad (12)$$

where  $G(h) = 12h(-144h^2 - 72bh + 3b^2 + 12b^3h - 16)$ ,

$$\begin{aligned} d_{00} &= \frac{3}{2}h, & d_{01} &= -\frac{1}{2} + \frac{3}{32}b^2, & d_{02} &= -\frac{1}{8}b, \\ d_{10} &= -\frac{1}{8}bh, & d_{11} &= h - \frac{9}{128}b^3 + \frac{3}{8}b, & d_{12} &= \frac{3}{32}b^2 - \frac{1}{3}, \\ d_{20} &= \frac{3}{128}b^2h - \frac{3}{8}h, & d_{21} &= \frac{27}{2048}b^4 - \frac{9}{64}b^2 + \frac{3}{8}, & d_{22} &= -\frac{9}{512}b^3 + \frac{5}{32}b + \frac{3}{4}h \end{aligned}$$

and

$$\begin{aligned} f_{00} &= 576h^2 + (-36b^3 + 192b)h, & f_{02} &= 12b^2 - 48bh - 64, \\ f_{10} &= -144bh^2 + (36b^2 - 192)h, & f_{12} &= 48h(b^2 - 4), \\ f_{20} &= (36b^2 - 576)h^2 + (27b^3 - 144b)h, & f_{22} &= 12h(-48h + 3b^3 - 16b). \end{aligned}$$

So, the integral  $I(h)$  can be expressed as the form

$$I(h) = \alpha(h)I_0 + (\beta(h) - \frac{3b}{4}\gamma(h))I_1 + \gamma(h)Z \quad (13)$$

and

$$I'(h) = \alpha_1(h)I'_0 + \beta_1(h)I'_1 + \gamma_1(h)Z' \quad (14)$$

where

$$\begin{aligned}
\alpha_1 &= \alpha' d_{00} + (\beta' - \frac{3}{4} b \gamma') d_{10} + \gamma' d_{20} + \alpha, \\
\beta_1 &= \alpha' d_{01} + (\beta' - \frac{3}{4} b \gamma') d_{11} + \gamma' d_{21} + \beta - \frac{3}{4} b \gamma, \\
\gamma_1 &= \alpha' d_{02} + (\beta' - \frac{3}{4} b \gamma') d_{12} + \gamma' d_{22} + \gamma
\end{aligned}$$

with  $\deg \alpha_1 \leq \left\lfloor \frac{n-1}{2} \right\rfloor$ ,  $\deg \beta_1 \leq \left\lfloor \frac{n-2}{2} \right\rfloor$ , and  $\deg \gamma_1 \leq \left\lfloor \frac{n-3}{2} \right\rfloor$ .

Let  $\Sigma = \{h | h > 0, \}$ ,  $\mathcal{N} = \{h | \beta_1(h) = 0 \text{ or } h = h_2, h \in \Sigma\}$ . Then, from [23] we obtain the following lemma.

**Lemma 4** For  $h \in \Sigma \setminus \mathcal{N}$

$$\left( \frac{I'(h)}{\beta_1(h)} \right)' = \frac{M(h)}{G(h)\beta_1^2(h)} \quad (15)$$

where

$$M(h) = \alpha_2(h)I'_0 + \beta_2(h)Z' \quad (16)$$

with  $\deg \alpha_2 \leq \left\lfloor \frac{n-1}{2} \right\rfloor + \left\lfloor \frac{n-2}{2} \right\rfloor + 2$ ,  $\deg \beta_2 \leq \left\lfloor \frac{n-1}{2} \right\rfloor + \left\lfloor \frac{n-2}{2} \right\rfloor + 1$ , and  $\alpha_2(0) = 0$ .

**Lemma 5** Let  $p$ ,  $q$ , and  $r$  be the number of zeros of  $I'(h)$ ,  $M(h)$ ,  $\beta_1(h)$  in the interval  $(0, +\infty)$ , respectively. Then

$$p \leq q + r + 1$$

**PROOF.** Let  $a_1, a_2, \dots, a_r$  be the zeros of  $\beta_1$  in the interval  $(0, +\infty)$  with  $0 = a_0 < a_1 < a_2 < \dots < a_r < +\infty$ , as shown in Fig. 2.




Fig. 2. The distribution of zeros of  $\beta_1$ .

Then, for each  $a_j, j = 1, \dots, r$ , there is a  $\delta > 0$  such that except  $a_j$  there are no zeros of  $I'$ ,  $M$  and  $\beta_1$  in the interval  $[a_j - \delta, a_j + \delta]$ .

Now, suppose  $a_j$  is the zero of  $I'$ ,  $M$  and  $\beta_1$  with multiplication  $l_j, m_j$ , and  $k_j$ , respectively. If  $k_j = l_j$ , then  $m_j - 2l_j \geq 0$ , which implies that  $m_j \geq l_j$ ; if  $k_j \neq l_j$ , then  $l_j - k_j - 1 = m_j - 2k_j$ , which also implies  $m_j \geq l_j$ .



On the other hand, let  $p_0, p_j$ , and  $p_r$  denote the number of zeros of  $I'$  in the intervals  $(a_0, a_1 - \delta)$ ,  $(a_j + \delta, a_{j+1} - \delta)$ , and  $(a_r + \delta, +\infty)$ , respectively, and let  $q_0, q_j$ , and  $q_r$  denote the number of zeros of  $M$  in the intervals  $(a_0, a_1 - \delta)$ ,  $(a_j + \delta, a_{j+1} - \delta)$ , and  $(a_r + \delta, +\infty)$ , respectively, where  $j = 1, \dots, r-1$ . Then, Eq. (15) implies  $p_j \leq q_j + 1, j = 0, \dots, r$ . Therefore,

$$p = \sum_{j=0}^r p_j + \sum_{j=1}^r l_j \leq \sum_{j=0}^r (q_j + 1) + \sum_{j=1}^r m_j = [q_0 + \sum_{j=1}^r (q_j + m_j)] + r + 1 = q + r + 1.$$

■

**Lemma 6** *Let  $W(h) = M(h)/I'_0$ . Then, we can obtain the Riccati equation*

$$G\beta_2 W' = R_2 + (G\beta'_2 + \beta_2(f_{22} - f_{00}) + 2\alpha_2 f_{02})W - f_{02}W^2 \quad (17)$$

*Furthermore, if letting  $s$  and  $t$  respectively denote the number of zeros of  $R_2$  and  $\beta_2$  in the interval  $(0, +\infty)$ , then we have*

$$p \leq r + s + t + 2$$

**PROOF.** Using interval  $(0, +\infty)$  instead of  $(0, h_2)$  in [23], the straightforward computing and Lemma 5 give these results.

■

Let  $\#\{I(h), h \in (m, n)\}$  denote the number of zeros of  $I(h)$  with respect to  $h$  in the interval  $(m, n)$ . Then, Eq. (14) implies

$$\#\{I(h), h \in (0, +\infty)\} \leq \#\{I'(h), h \in (0, +\infty)\} + 1$$

Since  $I(0) = R(0) = 0$  we have

$$\#\{I(h), h \in (0, +\infty)\} \leq \#\{I'(h), h \in (0, +\infty)\} + 1 - 1 \leq r + (s-1) + t + 2 \leq \frac{7n+3}{2}$$

**3.2 The case of global center:  $b > -2$  and  $3b^2 - 16 = 0$**

The equality  $3b^2 - 16 = 0$  means  $b = \frac{4\sqrt{3}}{3}$ .

In this section, we are going to use equations  $\mathbf{J}_2 = (I_0, Z)^T$ ,

$$I_1'' = \frac{\sqrt{3}}{3} I_0'' \quad (18)$$

and

$$G(h)\mathbf{J}_2'' = \begin{pmatrix} f_{00} & f_{02} \\ f_{20} & f_{22} \end{pmatrix} = \mathbf{F}_2\mathbf{J}_2' \quad (19)$$

instead of equations (10) and (12). We will also use equations (13) and (14) with  $b = \frac{4\sqrt{3}}{3}$ . Similar to the result given in section 3.1, the following result is obtained

$$\#\{I(h), h \in (0, +\infty)\} \leq \frac{7n+3}{2}$$

### 3.3 The case of double-cusps loop: $b = -2$ .

In this case,  $h = h_2 = \frac{1}{6}$  corresponds to the double-cusps loop. Since  $b = -2$ , equations (10) through (14) hold. Then, using  $\Sigma_1 = \{h|h > 0, h = h_2\}$  and  $\mathcal{N}_1 = \{h|\beta_1(h) = 0, h \in \Sigma_1\}$  instead of  $\Sigma$  and  $\mathcal{N}$ , we can obtain

$$\#\{I(h), h \in (0, +\infty)\} \leq \#\{I'(h), h \in (0, h_2)\} + \#\{I'(h), h \in (h_2, +\infty)\} \leq 7n+5$$

### 3.4 Main Result

The results given in subsections 3.1 through 3.3 are summarised in the following theorem.

**Theorem 1** *Suppose that  $I(h)$  is defined in Eq. (A) and  $a_{k,j} \in \mathbb{R}$ . The following relation holds*

$$\#\{I(h), h \in (0, +\infty)\} \leq 7n + 5.$$

## 4 Conclusions

In this paper, we have investigated the number of zeros of the Abelian integrals for a class of non-linear dynamical systems for the case  $b \geq -2$ . We have proved that the related elliptic integrals have at most  $7n + 5$  zeros including multiple zeros. Through the results presented in this paper and [23], we can obtain the upper bound of the number of limit cycles bifurcated from the periodical orbits for each  $b \in \mathbb{R}$ .

## References

- [1] Ananthkrishnan N., Sudershan S., Sudhakar K., Verma A., Large-amplitude limit cycles in resonantly coupled oscillators, *J. Sound Vibration*, 231 (2000) 1377–1382.
- [2] Arnold, V.I. “Ten Problems”. *Advance in Soviet Mathematics*. 1: 1-8 (1990).
- [3] Bejarano J. D., Garcia-Margallo J., The greatest number of limit cycles of the generalized Rayleigh-Linard oscillator, *J. Sound Vibration* 221 (1999) 133–142.
- [4] F. Dumortier, C. Rousseau, Cubic Liénard equations with linear damping, *Nonlinearity*, 3(1999), 1015-1039.
- [5] L.Gavrilov, The infinitesimal 16th Hilbert problem in the quadratic, *Invent. Math.* 143(2001)449-497.
- [6] Giacomini Hector, Neukirch Sbastien, Comment on “Linard systems, limit cycles, and Melnikov theory” *Phys. Rev. E* 59 (1999) 2483–2484.
- [7] Gueron Shay, Liron Nadav, Fast computation of limit cycles in an industrial application. Special issue on practical asymptotics, *J. Engrg. Math.*, 39 (2001) 79–86.
- [8] Han, M.. Cyclicity of planar homoclinic loops and quadratic integrable systems. *Science in China(ser:A)*. 40: 1247-1258 (1997).
- [9] Han, M.&Chen,J. The number of limit cycles in double homoclinic bifurcations. *Science in China(ser:A)*. 30: 401-414 (2000).
- [10] E.Horozov, I.D.Iliev, On the number of limit cycles in perturbations of quadratic Hamiltonian systems, *Proc. Lond. Math. Soc.*(3) 69(1994),198-224.
- [11] J.Li., Hilbert’s 16th problem and bifurcations of planar polynomial vector fields. *International Journal of Bifurcation and Chaos*. 13(1): 47-106 (2003).
- [12] Llibre Jaume, Pizarro Luis, Ponce Enrique, Limit cycles of polynomial Linard systems, *Phys. Rev. E*, 58 (1998) 5185–5187.
- [13] Matveev Alexey S., Savkin Andrey V., Limit cycles in a class of hybrid dynamical systems, *Math. Control Signals Systems* 15 (2002) 120–144.
- [14] Mickens Ronald E., Liénard systems, limit cycles, Melnikov theory, and the method of slowly varying amplitude and phase, *J. Sound Vibration* 217 (1998) 790–793.
- [15] G.S.Petrov, Number of zeros of complete elliptic integrals, *Funct. Anal. Anal.* 18(1984)72-73; 20(1986)37-40; 21(1987)87-88; 22(1988)37-40; 23(1989)88-89; 24(1990)45-50.
- [16] Sanjun Miguel A. F., Reply to “Comment on ‘Linard systems, limit cycles, and Melnikov theory’” *Phys. Rev. E* 59 (1999) 2485–2486.

- [17] Sez E., Sznt I., Monsalve A., Three limit cycles for a first-order exothermic chemical reaction in a continuous stirred-tank reactor, *J. Math. Chem.*, 27 (2000) 171–182.
- [18] S.Smale, “Dynamics restrospective: great problem, attempts that failed”. *Physica D.* 51: 261-273 (1991).
- [19] Wang Hsin-Hsiung, Krstić Miroslav, Extremum seeking for limit cycle minimization, *IEEE Trans. Automat. Control*, 45 (2000) 2432–2437.
- [20] T. Zhang, H. Zang, M. Han, Bifurcations of limit cycles in a cubic system. *Chaos Solitons Fractals* 20 (2004), no. 3, 629–638.
- [21] T. Zhang, M. Han, H. Zang, Perturbation from an asymmetric cubic Hamiltonian, *J. Math. Anal. Appl.* 305(2005) 617-630.
- [22] T. Zhang, M. O. Tade and Y. C. Tian, On the zeros of the Abelian integrals for a class of Liénard systems, *Physics Letters A* (submitted).
- [23] T. Zhang and H. Zang, On the number of zeros of Abelian integrals for a kind of Liénar system under perturbations, *Lond. Math. Soc.* (submitted).
- [24] Zhang W., Yu P., A study of the limit cycles associated with a generalized codimension-3 Lienard oscillator, *J. Sound Vibration*, 231 (2000) 145–173.
- [25] H. Zang, W. Chen, T. Zhang, Perturbation from a cubic Hamiltonian with three figure eight-loops. *Chaos Solitons Fractals* 22 (2004), no. 1, 61–74.
- [26] H. Zang, T. Zhang, W. Chen, Bifurcations of limit cycles from quintic Hamiltonian system with a double frgure eight loop, *J. Complexity*, 20 (2004) 544-560.